



thiamine-responsive megaloblastic anemia syndrome

Thiamine-responsive megaloblastic anemia syndrome is a rare condition characterized by hearing loss, diabetes, and a blood disorder called megaloblastic anemia.

Megaloblastic anemia occurs when a person has a low number of red blood cells (anemia), and the remaining red blood cells are larger than normal (megaloblastic).

The symptoms of this blood disorder may include decreased appetite, lack of energy, headaches, pale skin, diarrhea, and tingling or numbness in the hands and feet.

Individuals with thiamine-responsive megaloblastic anemia syndrome begin to show symptoms of megaloblastic anemia between infancy and adolescence. This syndrome is called "thiamine-responsive" because the anemia can be treated with high doses of vitamin B1 (thiamine).

People with thiamine-responsive megaloblastic anemia syndrome develop hearing loss caused by abnormalities of the inner ear (sensorineural hearing loss) during early childhood. It remains unclear whether thiamine treatment can improve hearing or prevent hearing loss.

Diabetes becomes apparent in affected individuals sometime between infancy and adolescence. Although these individuals develop diabetes during childhood, they do not have the form of the disease that develops most often in children, called type 1 (autoimmune) diabetes. People with thiamine-responsive megaloblastic anemia syndrome usually require insulin to treat their diabetes. In some cases, treatment with thiamine can reduce the amount of insulin a person needs.

Some individuals with thiamine-responsive megaloblastic anemia syndrome develop optic atrophy, which is the degeneration (atrophy) of the nerves that carry information from the eyes to the brain. Heart and blood vessel (cardiovascular) problems such as heart rhythm abnormalities and heart defects have also been reported in some people with this syndrome.

Frequency

Thiamine-responsive megaloblastic anemia syndrome has been reported in approximately 30 families worldwide. Its prevalence is unknown.

Genetic Changes

Mutations in the *SLC19A2* gene cause thiamine-responsive megaloblastic anemia syndrome. This gene provides instructions for making a protein called thiamine transporter 1, which transports thiamine into cells. Thiamine is found in many different foods and is important for numerous body functions.

Most mutations in the *SLC19A2* gene lead to the production of an abnormally short, nonfunctional thiamine transporter 1. Other mutations change single protein building blocks (amino acids) in this protein. All of these mutations prevent thiamine transporter 1 from bringing thiamine into the cell. It remains unclear how the absence of this protein leads to the seemingly unrelated symptoms of megaloblastic anemia, diabetes, and hearing loss. Research suggests that an alternative method for transporting thiamine is present in all the cells of the body, except where blood cells and insulin are formed (in the bone marrow and pancreas, respectively) and cells in the inner ear.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- Rogers syndrome
- Thiamine-responsive myelodysplasia
- TRMA

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Megaloblastic anemia, thiamine-responsive, with diabetes mellitus and sensorineural deafness
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0342287/>

Other Diagnosis and Management Resources

- GeneReview: Thiamine-Responsive Megaloblastic Anemia Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1282>
- MedlinePlus Encyclopedia: Optic nerve atrophy
<https://medlineplus.gov/ency/article/001622.htm>
- MedlinePlus Encyclopedia: Thiamine
<https://medlineplus.gov/ency/article/002401.htm>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>

- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Megaloblastic Anemia (image)
<https://medlineplus.gov/ency/imagepages/1214.htm>
- Encyclopedia: Optic nerve atrophy
<https://medlineplus.gov/ency/article/001622.htm>
- Encyclopedia: Thiamine
<https://medlineplus.gov/ency/article/002401.htm>
- Health Topic: Anemia
<https://medlineplus.gov/anemia.html>
- Health Topic: Diabetes
<https://medlineplus.gov/diabetes.html>
- Health Topic: Hearing Problems in Children
<https://medlineplus.gov/hearingproblemsinchildren.html>

Genetic and Rare Diseases Information Center

- Thiamine responsive megaloblastic anemia syndrome
<https://rarediseases.info.nih.gov/diseases/9210/thiamine-responsive-megaloblastic-anemia-syndrome>

Educational Resources

- Disease InfoSearch: Thiamine responsive megaloblastic anemia syndrome
<http://www.diseaseinfosearch.org/Thiamine+responsive+megaloblastic+anemia+syndrome/7055>
- Kids Health: Diabetes Center
<http://kidshealth.org/en/parents/center/diabetes-center.html>
- MalaCards: thiamine-responsive megaloblastic anemia syndrome
http://www.malacards.org/card/thiamine_responsive_megaloblastic_anemia_syndrome

- March of Dimes: Hearing Impairment
<http://www.marchofdimes.org/baby/hearing-impairment.aspx>
- Orphanet: Thiamine-responsive megaloblastic anemia syndrome
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=49827

Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD): Megaloblastic Anemia
<https://rarediseases.org/rare-diseases/anemia-megaloblastic/>
- Resource list from the University of Kansas Medical Center: Diabetes
<http://www.kumc.edu/gec/support/diabetes.html>
- Resource list from the University of Kansas Medical Center: Hard of Hearing/Deafness
<http://www.kumc.edu/gec/support/hearing.html>

GeneReviews

- Thiamine-Responsive Megaloblastic Anemia Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1282>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Anemia,+Megaloblastic%5BM%5D%29+AND+%28%28thiamine-responsive+megaloblastic+anemia+syndrome%5BTIAB%5D%29+OR+%28thiamine-responsive%5BTIAB%5D+AND+megaloblastic+anemia%5BTIAB%5D+AND+syndrome%5BTIAB%5D%29+OR+%28rogers+syndrome%5BTIAB%5D%29+OR+%28trma%5BTIAB%5D%29+OR+%28thiamine-responsive+myelodysplasia%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D%5D>

OMIM

- THIAMINE-RESPONSIVE MEGALOBlastic ANEMIA SYNDROME
<http://omim.org/entry/249270>

Sources for This Summary

- Baron D, Assaraf YG, Cohen N, Aronheim A. Lack of plasma membrane targeting of a G172D mutant thiamine transporter derived from Rogers syndrome family. *Mol Med*. 2002 Aug;8(8):462-74.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12435857>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2040007/>
- Diaz GA, Banikazemi M, Oishi K, Desnick RJ, Gelb BD. Mutations in a new gene encoding a thiamine transporter cause thiamine-responsive megaloblastic anaemia syndrome. *Nat Genet*. 1999 Jul;22(3):309-12.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10391223>

- GeneReview: Thiamine-Responsive Megaloblastic Anemia Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1282>
- Liberman MC, Tartaglioni E, Fleming JC, Neufeld EJ. Deletion of SLC19A2, the high affinity thiamine transporter, causes selective inner hair cell loss and an auditory neuropathy phenotype. *J Assoc Res Otolaryngol*. 2006 Sep;7(3):211-7. Epub 2006 Apr 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16642288>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1805778/>
- Lorber A, Gazit AZ, Khoury A, Schwartz Y, Mandel H. Cardiac manifestations in thiamine-responsive megaloblastic anemia syndrome. *Pediatr Cardiol*. 2003 Sep-Oct;24(5):476-81. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14627317>
- Olsen BS, Hahnemann JM, Schwartz M, Østergaard E. Thiamine-responsive megaloblastic anaemia: a cause of syndromic diabetes in childhood. *Pediatr Diabetes*. 2007 Aug;8(4):239-41.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17659067>
- Ricketts CJ, Minton JA, Samuel J, Ariyawansa I, Wales JK, Lo IF, Barrett TG. Thiamine-responsive megaloblastic anaemia syndrome: long-term follow-up and mutation analysis of seven families. *Acta Paediatr*. 2006 Jan;95(1):99-104.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16373304>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/thiamine-responsive-megaloblastic-anemia-syndrome>

Reviewed: February 2009

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services